3 (Sem-1/CBCS) MAT HC 1

2021 (Held in 2022)

MATHEMATICS

(Honours)

Paper: MAT-HC-1016

(Calculus)

Full Marks: 60


Time: Three hours

The figures in the margin indicate full marks for the questions.

- 1. Answer the following questions: 1×7=7
 - (a) Write down the nth derivative of $y = \log x$.
 - (b) The point P(c,f(c)) on the graph of f(x) is such that f'(c)=0. Does it necessarily imply that P is an inflection point on the graph?

- (c) Write down the value of $\lim_{x \to \infty} x \sin \frac{1}{x}$.
- (d) Find the domain of the vector function $\vec{F}(t) = (1-t)\hat{i} + \sqrt{-t} \hat{j} + \frac{1}{t-2}\hat{k}$
- (e) Write one basic difference between the disk/washer and shell method for computing volume of revolution.
- (f) What is the direction of velocity of a moving object on its trajectory.
- (g) The velocity of a particle moving in space is $\vec{V}(t) = e^t \hat{i} + t^2 \hat{j}$. Find the direction of motion at time t = 2.
- 2. Answer the following questions: 2×4=8
 - (a) Applying L. Hopital's rule, evaluate $\lim_{x\to \frac{\pi}{4}} (1 \tan x) \sec 2x$
 - (b) Write down the parametric equation of a line that contains the point (3,1,4) and is parallel to the vector $\vec{v} = -\hat{i} + \hat{j} 2\hat{k}$.
 - (c) Find the area of the surface generated by revolving the portion of the curve $y = x^3$ between x = 0 and x = 1 about the x-axis.

- (d) Explain briefly why the acceleration of an object moving with constant speed is always orthogonal to the direction of motion.
- 3. Answer any three of the following questions: 5×3=15
 - (a) If $y = \cos(m \sin^{-1} x)$, show that $(1-x^2)y_{n+2} (2n+1)x y_{n+1} + (m^2 n^2)y_n = 0.$ Hence find $y_n(0)$. 3+2=5
 - (b) Sketch the graph of a function f with all the following properties: 5
 - (i) the graph has y=1 and x=3 as asymptotes
 - (ii) f is increasing for x < 3 and 3 < x < 5 and decreasing elsewhere
 - (iii) the graph is concave up for x < 3 and concave down for 3 < x < 7
 - (iv) f(0) = 4 = f(5) and f(7) = 2



(d) Obtain the reduction formula for $\int tan^n x \, dx$.

Hence evaluate $\int_{0}^{\pi/4} \tan^{5} x \, dx \qquad 3+2=5$

The position vector of a moving object at any time t is given by $R(t) = ti + e^{t}j$. Find the tangential and normal components of the object's acceleration.

4. Answer any three of the following questions: 10×3=30

(a) A firm determines that x units of its product can be sold daily at rupees p per unit where x=1000-p. The cost of producing x units per day is

C(x) = 3000 + 20x. Then —

(i) Find the revenue function R(x).

(ii) Find the profit function p(x). 2

- is atmost 500 units per day, determine how many units the company must produce and sell each day to maximize profit. 3
- (iv) Find the maximum profit. 2
- (v) What price per unit must be charged to obtain maximum profit?
- (b) When is an object said to move in central force field? Derive Kepler's 2nd law of motion, assuming that planetary motion occurs in central force field.

 2+8=10
- (c) (i) Find the length of the arc of the astroid $x^{2/3} + y^{2/3} = 1$ lying in the positive quadrant.
 - (ii) Using cylindrical shell method, find the volume of the solid formed by revolving the region bounded by the parabola $y = 1 x^2$, the y-axis, and the positive x-axis, about y-axis.

(in) Find the surface area generated when the polar curve

r=5, 0505%

is revolved about x-axis.

3

- (d) (i) Find the volume generated by disk/washer method, when the region bounded by y = x, y = 2x and y = 1 is revolved about the x-axis
 - (ii) A particle moves along the polar path (r, θ) where

 $r(t) = 3 + 2\sin t, \theta(t) = t^3.$

Find the velocity $\vec{v}(t)$ and acceleration $\vec{A}(t)$ in terms \hat{u}_t and \hat{u}_{θ} .

- (i) Evaluate $\lim_{x\to 0} (1 + \sin x)^{1/x}$. 3
 - (ii) Examine the existence of vertical tangent and cusp of the graph of $y = (x-4)^{3/2}$.
 - (iii) A projectile is fired from ground level at an angle of 30° with muzzle speed 110 ft/sec. Find the time of flight and the range.

(i) Obtain the reduction formula for $\int \cos^n x \, dx$.

Hence evaluate $\int \cos^5 x \, dx$.

3+2=5

(ii) Find the unit tangent vector $\vec{T}(t)$ and principal unit normal vector $\vec{N}(t)$ at each point on the graph of vector function

 $\vec{R}(t) = (3\sin t, 4t, 3\cos t)$ 5

3 (Sem-1/CBCS) MAT HC2

2021

(Held in 2022)

MATHEMATICS

(Honours)

Paper: MAT-HC-1026

(Algebra)

Full Marks: 80

Time: Three hours

The figures in the margin indicate full marks for the questions.

1. Answer the following as directed:

1×10=10

- (a) Find the polar representation of z = 2i.
- (b) If x = 0 and y > 0, then what is the value of t^* ?
- (c) Write the negation of the statement 'For any integer n, n² > n' in plain English then formulate the negation using set of context and quantifier.

- (d) Disapprove the statement using counter example:
 - "For any $x, y \in \mathbb{R}$, $x^2 = y^2$ implies x = y."
- (e) Suppose f is a constant function from X to Y. The inverse image of a subset of Y cannot be
 - (i) an empty set
 - (ii) the whole set X
 - (iii) a non-empty proper subset of X (Choose the correct option)
- (f) Let $X = Y = \mathbb{R}$. Let $A \subseteq X$, $B \subseteq Y$. Draw the picture for $A \times B$ where A = [-1,1] and B = [2,3].
- (g) Suppose a system of linear equations in echelon form has a 3 × 5 augmented matrix whose fifth column is a pivot column.
 Is the system consistent? Justify.
- (h) If a set $S = \{\bar{v}_1, \bar{v}_2, ..., \bar{v}_p\}$ in \mathbb{R}^n contains the \bar{O} vector, is the set linearly independent? Justify.

(i) If
$$A = \begin{bmatrix} 1 & -3 \\ -2 & 4 \end{bmatrix}$$
 $\hat{x} = \begin{bmatrix} 5 \\ 3 \end{bmatrix}$, compute $(A\hat{x})^T$

- What is the determinant of an n×n elementary matrix E that has been scaled by 7.
- 2. Answer the following questions: 2×5=10
 - (a) If $z = -2\sqrt{3} 2i$, find the polar radius and polar argument of z.
 - (b) Is the function $g: \mathbb{R} \to \mathbb{R}$ given by g(x) = |x-2| one-one and onto? Explain.
 - (c) Let universal set be \mathbb{R} and index set be \mathbb{R} . For a natural number n, $J_n = \left(0, \frac{1}{n}\right)$.

Identify with justification $\bigcap_{n\in N} J_n$.

(d) Show that T is a linear transformation by finding a matrix that implements the mapping

$$T(x_1, x_2, x_3, x_4) = (0, x_1 + x_2, x_2 + x_3, x_3 + x_4)$$

- (c) A is a 2 × 4 matrix with two pivot positions. Answer the following with justification:
 - (i) Does $A\vec{x} = \vec{0}$ have a non-trivial solution?
 - (ii) Does $A\vec{x} = \vec{b}$ have at least one solution for every \vec{b} ?
- 3. Answer any four questions from the following: 5×4=20
 - (a) Find the polar representation of the complex number
 z=1-cosa+isina a∈ [0, 2π).
 - (b) Let A and B be subsets of an universal set U. Prove —

(i)
$$(A \cap B)^C = A^C \cup B^C$$

(ii)
$$(A \cup B)^C = A^C \cap B^C$$

- (c) Define bijection.

 Let $f: \mathbb{N} \to \mathbb{N}$ be -f(m) = m-1. If m is even f(m) = m+1, if m is odd. Show f is a bijection and $f^{-1} = f$. 1+4=5
- (d) For vectors $\vec{v}_1, \vec{v}_2, ..., \vec{v}_p \in \mathbb{R}^n$ define span $\{\vec{v}_1, \vec{v}_2, ..., \vec{v}_p\}$ construct a 3×3 matrix A with non-zero elements and a vector \vec{b} on \mathbb{R}^3 such that \vec{b} is not in the set spanned by the columns of A. 2+3=5
- (e) Alka-Seltzer contains sodium bicarbonate (NaHCO₃) and citric acid (H₃C₆H₅O₇). When a tablet is dissolved in water the following reaction produces sodium citrate, water and carbon dioxide:
- $NaHCO_3 + H_3C_6H_5O_7 \rightarrow Na_3C_6H_5O_7 + H_2O + CO_2$ Balance the chemical equation using vector equation approach.
 - (f) Prove that an n × n matrix A is invertible if and only if A is row equivalent to Invand in this case any sequence of elementary row operations that reduces A to In also transforms In into A-1.

5

4. Answer any four from the following: 10×4=40

- (a) (i) Find the cube roots of the number z = 1 + i and represent them in the complex plane.
 - (ii) Find the number of ordered pairs (a, b) of real numbers such that $(a+ib)^{2002} = a-ib$.
 - (iii) If x, y, z be real numbers such that $\sin x + \sin y + \sin z = 0$ and $\cos x + \cos y + \cos z = 0$, prove that $\sin 2x + \sin 2y + \sin 2z = 0$ and $\cos 2x + \cos 2y + \cos 2z = 0$.

3

(b) (i) Solve the equation $z^7 - 2iz^4 - iz^3 - 2 = 0.$ 5

(ii) Find the inverse of the matrix if it exists by performing suitable row operations on the augmented matrix [A:I]

$$A = \begin{bmatrix} 1 & 0 & -2 \\ -3 & 1 & 4 \\ 2 & -3 & 4 \end{bmatrix}$$
 5

- (i) If $f: X \to Y$ be a map and $B \subseteq Y$, then prove $f^{-1}(B^C) = (f^{-1}(B))^C$.
 - (ii) $A_n = \left(-\frac{1}{n}, \frac{1}{n}\right)$, where $n \in \mathbb{N}$. Find $\bigcup A_n$ and $\bigcap A_n$.
 - (iii) Let $f:\mathbb{R} \to \mathbb{R}$ be given $f(x) = x^2$.

Find $f^{-1}(1)$, $f^{-1}(-1)$, $f^{-1}([0, 1])$.

State the induction principle and (d) (i) use it to show that for any positive integer $1+2+3+...+n=\frac{n(n+1)}{2}$.

4

4

2

Write as an implication 'square of (ii) an even number is divisible by 4'. Then use direct proof to prove it.

- (iii) Give proof using contrapositive 'For an integer x if x^2-6x+5 is even, then x is odd'.
- (e) (i) Use the invertible matrix theorem to decide if A is invertible

$$A = \begin{bmatrix} 1 & 0 & 2 \\ 3 & 1 & 2 \\ -5 & -1 & 9 \end{bmatrix}$$

(ii) Compute det A where

$$A = \begin{bmatrix} 2 - 8 & 6 & 8 \\ 3 - 9 & 5 & 10 \\ -3 & 0 & 1 - 2 \\ 1 - 4 & 0 & 6 \end{bmatrix}$$

(iii) What do you mean by equivalence class for an equivalence relation?
 For the relation a = b mod(5) on z, find all the distinct equivalence classes of z.
 1+3=4

(i) Solve the system of equations

$$x_1 - 3x_3 = 8$$

$$2x_1 + 2x_2 + 9x_3 = 7$$

$$x_2 + 5x_3 = -2$$

4

- (ii) Choose h and k such that the system has 4
 - (a) no solution
 - (b) a unique solution
 - (c) many solutions $x_1 + hx_2 = 2$ $4x_1 + 8x_2 = k$
- (iii) Write the general solution of $10x_1 3x_2 2x_3 = 7$ in parametric vector form.
- (g) (i) Prove that the indexed set $S = \{\vec{v}_1, \vec{v}_2, ..., \vec{v}_p\}$ of two or more vectors is linearly dependent if and only if at least one of the vectors in S is a linear combination of the others. In fact, if S is linearly dependent and $\vec{v}_1 \neq \vec{0}$, then some \vec{v}_j (with j > 1) is a linear combination of the preceding vectors $\vec{v}_1, \vec{v}_2, ..., \vec{v}_{j-1}$.

(ii) Use Cramer's rule to compute the solutions of the system 3

$$-5x_1 + 3x_2 = 9$$
$$3x_1 - x_2 = -5$$

- (iii) Suppose $T: \mathbb{R}^5 \to \mathbb{R}^2$ and $T(\bar{x}) = A\bar{x}$ for some matrix A and each \bar{x} in \mathbb{R}^5 . How many rows and columns does A have? Justify.
- (h) (i) Let T: ℝ² → ℝ² be the transformation that rotates each point in ℝ² about the origin through an angle φ with the counter-clockwise direction taken as positive. Find the standard matrix for this transformation.

(ii) Let $T: \mathbb{R}^n \to \mathbb{R}^m$ be a linear transformation. Prove that T is one-to-one if and only if the equation $T(\bar{x}) = \bar{0}$ has only the trivial solution.

3

(iii) Find the area of the parallelogram whose vertices are (0, -2), (6, -1), (-3, 1) and (3, 2).